

Albano and Saracco reply. In our paper [1], it is assumed that, at criticality and starting from a ground-state configuration, the order parameter (OP) decays according to

$$OP(t) \propto t^{-\beta/\nu_{\perp}z}, \quad (1)$$

(Eq. (9) in [1]), where it is implicit that $z = z_{\parallel}$. In the preceding comment, Caracciolo et al. (CGGP) [2] have suggested, instead, that z must be interpreted as $z_{\perp}$. Based on this assumption CGGP conclude that the exponent $\beta/\nu_{\perp}z$ reported in [1] supports the field-theoretical equation developed in reference [3] (Eq.(1) in [1]), while the numerical estimate of the exponent $c_{\perp}$ differs from the prediction of both field-theoretical models considered by us, given by Eqs.(1) and (2) in [1], which were taken from references [3] and [4], respectively. The conclusions of CGGP are in contrast to the main finding of our paper that fully supports the universality class predicted by Eq.(2) in [1].

Let us now show that the agreement between the exponent $\beta/\nu_{\perp}z_{\perp}$ predicted by Eq.(1) in [1] and our numerical data is merely coincidental and that the conclusions of CGGP are inconsistent since the relevance of the exponent $c_{\perp}$ to determine the universality class is simply disregarded.

Taking the logarithmic derivative of Eq.(1) at criticality one has

$$\partial \ln OP(t, \tau)|_{\tau=0} \equiv OP^* \propto t^{1/\nu_{\perp}z}. \quad (2)$$

Furthermore, inserting Eq.(2) in Eq.(1) one gets $\ln(OP) \propto -\beta \ln(OP^*)$, which gives an unbiased estimation of β independent of any assumption about z . Our data shown in Figure 1 are well fitted by an exponent $\beta = 0.330$, in agreement with the universality class predicted by Eq.(2) in [1], while the value $\beta = 1/2$ predicted by Eq.(1) in [1] can clearly be ruled out.




FIG. 1. Log-log plot of OP versus the logarithmic derivative of OP . The full line with slope $\beta = -0.330$ is the best fit of the data. The dashed line has slope $\beta = 1/2$.

From Eq.(2) it also follows that $OP^{\nu_{\perp}} \propto t^{1/z}$. Our numerical results shown in figure 2 are in full agreement with $z = z_{\parallel} \simeq 1.998$, while the value $z_{\perp} \simeq 4$ suggested by CGGP can clearly be ruled out.




FIG. 2. Log-log plots of the $OP^{\nu_{\perp}}$ versus t . The full line with slope $1/z_{\parallel} = 0.501$ is the best fit of the data. The dashed line has slope $1/z_{\perp} = 0.25$.

Summing up, we have provide conclusive evidence showing that, in contrast to the suggestions of the preceding comment [2]: 1) The decay of the order parameter is governed by the time evolution of the correlation length parallel to the field and consequently the exponent z of Eq.(1) must be identified as $z_{\parallel}$ as in our original paper [1]; 2) A numerical determination of the exponent β can be performed independently of any assumption on the value of z ; 3) All exponents measured are fully consistent with the universality class predicted by Eq.(2) in [1] and developed in reference [4].

Acknowledgments: This work was supported by CONICET, UNLP and ANPCyT (Argentina). G.S. acknowledges the CIC for a research fellowship.

Ezequiel V. Albano and Gustavo Saracco. Instituto de Investigaciones Fisicoquímicas Teóricas y Aplicadas (INIFTA), UNLP, CONICET, Casilla de Correo, 16 Sucursal 4, (1900) La Plata, Argentina.

-
- [1] E. V. Albano and G. Saracco. Phys. Rev. Lett, **88** 145701 (2002).
 - [2] S. Caracciolo, A. Gambassi, M. Gubinelli and A. Pelissetto, preceeding comment, Phys. Rev. Lett. **XX**, xxxxxx (2003).
 - [3] H. K. Janssen and B. Schmittmann, Z. Phys. B **64**. 503 (1986). K.-t. Leung and J. L. Cardy, J. Stat. Phys. **44**, 567 (1986); ibid **45**, 1087 (Erratum) (1986).
 - [4] P. L. Garrido, F. de los Santos and M. A. Muñoz, Phys. Rev. E **57**, 752 (1998), and P. L. Garrido, M. A. Muñoz, and F. de los Santos, Phys. Rev. E **61**, R4683 (2000).